

Improving Patient Medication Adherence: A \$290 Billion Opportunity



Reducing the \$290 billion spent annually because of poor medication adherence requires building on proven practices and implementing policy actions that target the root causes of the problem.²

Poor medication adherence represents a significant source of wasteful health care spending. The causes of non-adherence are complex and systemic, resulting from high out-of-pocket costs, poor care coordination and the failure to account for the patient's personal circumstances.

Improving medication adherence requires building on a coordinated set of proven practices in the field and policy actions in both the public and private sectors.

THE PROBLEM

Scope of Poor Medication Adherence

- Of the approximately 187 million Americans who take one or more prescription drugs, up to one-half do not take their medications as prescribed.^{3,4}
- As many as 2 billion cases of poor medication adherence each year are avoidable.^{5,6}

Costs of Poor Medication Adherence

- Not taking medications as prescribed costs over \$100 billion a year in excess hospitalizations.⁷
- Total annual health care spending for a diabetes patient with low medication adherence (\$16,499) is almost twice the amount for a patient with high adherence (\$8,886).⁸
- Among hypertension patients, an estimated 89,000 premature deaths per year could be avoided with appropriate medication treatment.⁹
- Diabetes patients with poor medication adherence have a 30 percent yearly risk of hospitalization, as opposed to a 13 percent risk for those who accurately follow prescriber guidelines.¹⁰
- Non-adherent diabetes and heart disease patients have significantly higher mortality rates (12.1 percent) than similar patients who were adherent (6.7 percent).¹¹

Causes of Poor Medication Adherence¹²

- High out-of-pocket costs, especially for patients on multiple prescriptions for chronic conditions.
- Lack of care coordination, follow-up and shared decision-making.
- Complex or burdensome treatment regimens or multiple prescribed medications.
- Co-morbidities, such as severe and persistent mental illness.
- Side effects of prescribed medications, whether real or perceived.
- Personal factors, including lifestyle, culture and belief system.

SOLUTIONS

Improve Care Coordination

- **Proven Practice:** Care teams composed of physicians, pharmacists, nurses and other health care professionals can more effectively monitor adherence and counsel patients.¹³
- **Proven Practice:** Diabetes patients receiving case management, including bi-weekly automated calls and self-care training by nurses, are 21 percent more adherent to their medications than those who receive usual care.¹⁴

Enhance Patient Engagement and Education

- **Proven Practice:** Elderly patients who receive pharmacist-led discharge counseling before

Continued on back

Using care coordination strategies, patient engagement and Medication Therapy Management can significantly improve medication adherence.

Improving medication adherence also requires investments in HIT and financial incentives for patients and providers.

hospital discharge improve their medication adherence by 43 percent.¹⁵

- **Proven Practice:** Patients who participate in motivational interviewing and discussions about their individual needs, constraints and preferences are 13 percent more likely to take their medications as prescribed compared to patients receiving usual care.¹⁶
- **Proven Practice:** Patients with depression who are provided educational materials and one-on-one follow-up are twice as likely to refill their prescriptions.¹⁷

Utilize Counseling and Medication Management

- **Proven Practice:** Fifty-six percent of HIV/AIDS patients enrolled in a Medication Therapy Management (MTM) program, a multi-disciplinary team approach to care, follow their medication directions, as compared to 38 percent of patients who did not receive MTM.¹⁸
- **Proven Practice:** Patients with high blood pressure taking once-daily therapies are 11 percent more adherent than those taking twice-daily therapies.¹⁹

Expand Screening and Assessment

- **Proven Practice:** Expanding the use of proven screening and assessment tools to target patients at greatest risk for non-adherence, such as those with depression.²⁰
- **Proven Practice:** Establishing tools for providers to promote medication review and reconciliation as well as patient engagement, such as the American Society of Health-System Pharmacists Medication Reconciliation Toolkit.²¹

Invest in HIT Infrastructure²²

- **Policy Action:** Invest in electronic health records, e-Prescribing, clinical decision support systems and sharing of data related to the proper use of medications.
- **Policy Action:** Encourage sharing of near real-time prescription fill and refill data among providers, between patients and providers, and between providers and pharmacists to implement instantaneous point-of-care medication review and regimen reconciliation.

Employ Quality Measurement

- **Policy Action:** Adopt consensus-based standards, such as those from the National Quality Forum and Pharmacy Quality Alliance, to measure the quality of adherence strategies.^{23,24,25}
- **Policy Action:** Develop specific measures for adherence to medications for chronic disease.

Establish Financial Incentives²⁶

- **Policy Action:** Provide incentives for Medication Therapy Management and patient counseling.
- **Policy Action:** Eliminate co-payments for generic drugs and reduce brand-name co-payments.²⁷
- **Policy Action:** Expand adoption of value-based insurance design to reduce co-payments for medications for chronic conditions.
- **Policy Action:** Enable prescribers to simplify dosing by considering adherence and simplification of medication regimens in the development of formularies and cost-sharing requirements.

► Learn more about ways to Bend the Curve in health care costs at: www.nehi.net/bendthecurve

THE PROBLEM

1. NEHI. (2008). How Many More Studies Will It Take? A Collection of Evidence That Our Health Care System Can Do Better. Retrieved from http://www.nehi.net/publications/30/how_many_more_studies_will_it_take. Last accessed October 2011.
2. NEHI. (2009). Thinking Outside the Pillbox: A System-wide Approach to Improving Patient Medication Adherence for Chronic Disease. Retrieved from http://www.nehi.net/publications/44/thinking_outside_the_pillbox_a_systemwide_approach_to_improving_patient_medication_adherence_for_chronic_disease. Last accessed October 2011.
3. Kaiser Family Foundation. Prescription Drug Trends, May 2010. Retrieved from <http://www.kff.org/rxdrugs/upload/3057-08.pdf>. Last accessed October 2011.
4. Osterberg, L., Blaschke, T. (2005). Adherence to medication. *N Engl J Med*, 353(5), 487-497.
5. Osterberg and Blaschke. 2005.
6. IMS Health. (2010). National Prescription Audit PLUS. Retrieved from http://www.imshealth.com/deployedfiles/imshealth/Global/Content/StaticFile/Top_Line_Data/2010_Top_Therapeutic_Classes_by_RX.pdf. Last accessed October 2011.
7. Sokol, M.C., McGuigan, K.A., Verbrugge, R.R., et al. (2005). Impact of medication adherence on hospitalization risk and healthcare cost. *Med Care*, 43(6), 521-530.
8. Ho, P.M., Magid, D.J., Masoudi, F.A., et al. (2006). Adherence to cardioprotective medications and mortality among patients with diabetes and ischemic heart disease. *BMC Cardiovasc Disord*, 6, 48.

9. Cutler, D.M., Long, G., Berndt, E.R., et al. (2007). The value of antihypertensive drugs: A perspective on medical innovation. *Health Aff*, 26(1), 97-110.
10. Sokol, McGuigan, and Verbrugge. 2005.
11. Ho, Magid, Masoudi, et al. 2006.
12. Osterberg and Blaschke. 2005.

SOLUTIONS

13. NEHI. (2010). Thinking Outside the Pillbox, Medication Adherence and Care Teams: A Call for Demonstration Projects. Retrieved from http://www.nehi.net/publications/48/medication_adherence_and_care_teams_a_call_for_demonstration_projects. Last accessed October 2011.
14. Piette, J.D., Weinberger, M., McPhee, S.J., et al. (2000). Do automated calls with nurse follow-up improve self-care and glycemic control among vulnerable patients with diabetes? *Am J Med*, 108(1), 20-27.
15. Lipton, H.L., Bird, J.A. (1994). The impact of clinical pharmacists' consultations on geriatric patients' compliance and medical care use: A randomized controlled trial. *Gerontologist*, 34(3), 307-315.
16. Ogedegbe, G., Chaplin, W., Schoenthaler, A., et al. (2008). A practice-based trial of motivational interviewing and adherence in hypertensive African Americans. *Am J Hypertens*, 21(10), 1137-1143.
17. Katon, W., Rutter, C., Ludman, E.J., et al. (2001). A randomized trial of relapse prevention of depression in primary care. *Arch Gen Psychiatry*, 58(3), 241-247.
18. Hirsch, J.D., Rosenquist, A., Best, B.M., et al. (2009). Evaluation of the first year of a pilot program in community pharmacy: HIV/AIDS medication therapy management for

- Medi-Cal beneficiaries. *J Manage Care Pharm*, 15(1), 32-41.
19. Mounier-Vehier, C., Bernaud, C., Carre, A., et al. (1998). Compliance and antihypertensive efficacy of amlodipine compared with nifedipine slow-release. *Am J Hypertens*, 11(4 Pt 1), 478-486.
20. American Association of Colleges of Pharmacy. (2009). Better Medication Adherence is Essential to Improve Health Care Quality, Outcomes and Value. Alexandria, VA. Retrieved from <http://www.aacp.org/issuesandadvocacy/advocacy/SignonLetters/Documents/Policy%20Recommendations%2010-14-09.pdf>. Last accessed October 2011.
21. American Society of Health-System Pharmacists. (2011). ASHP Medication Reconciliation Toolkit, Bethesda, MD. Retrieved from http://www.ashp.org/Import/PRACTICEANDPOLICY/PracticeResourceCenters/PatientSafety/ASHP-MedicationReconciliationToolkit_1.aspx. Last accessed October 2011.
22. American Association of Colleges of Pharmacy. 2011.
23. American Association of Colleges of Pharmacy. 2011.
24. National Quality Forum. (2010). National Voluntary Consensus Standards for Medication Management. Washington, DC.
25. Pharmacy Quality Alliance. (2010). Pharmacy Quality Alliance Approved Measures, Washington, DC. Available at: <http://www.pqaalliance.org/files/PQA%20approved%20measures.pdf>. Last accessed October 2010.
26. American Association of Colleges of Pharmacy. 2011.
27. Maciejewski ML, Farley JF, Parker J, et al. Copayment reductions generate greater medication adherence in targeted patients. *Health Aff*, 2010;29(11):2002-2008.